VACCINES GIVEN TO **NEW BRUNSWICK CHILDREN** | PRENATAL-18 YEARS



4 DOSES of Diphtheria 4 DOSES of Tetanus 4 DOSES of Pertussis 1 DOSE of Polio (1957)

1 DOSE of Smallpox

1983 22 DOSES of 7 Vaccines



3 DOSES of Pneumococcal C-13 3 DOSES of Hepatitis B 3 DOSES of Rotavirus 2 DOSES of Measles 2 DOSES of Mumps

2019 70 DOSES of 16 Vaccines



2 DOSES of Rubella 2 DOSES of Varicella (Chickenpox) 2 DOSES of HPV 1 DOSE of Meningococcal C-C 1 DOSE of Meningococcal C-ACYW 7 DOSES of Diphtheria 7 DOSES of Tetanus 7 DOSES of Pertussis 5 DOSES of Polio (IPV) 4 DOSES of Hib

THE CHOICE IS YOURS

In New Brunswick parents still have the right to refuse any or all vaccines for infants & children. For children to attend school or day care a simple letter of refusal is sufficient. You also have the right to refuse disclosure of your child's private medical information. Health Canada has stated: "Unlike some countries, immunization is not mandatory in Canada, it cannot be made mandatory because of the Canadian Constitution...legislation and regulations must not be interpreted to imply compulsory immunization." Visit our website for more information.

What's in those vaccines?

- Complex biochemical substances including DNA debris from animal and aborted fetal tissue, foreign proteins, cell fragments, bovine serum albumin, viral and bacterial particles, emulsifiers, neurotoxins, formaldehyde, monosodium L-glutamate, egg proteins and much more.
- Aluminum Adjuvant: A neurotoxin that triggers brain inflammation and is linked to brain injuries and autoimmune disorders. 7210 micrograms injected in first year.
- Mercury Preservative: A neurotoxin contained in some multi-dose vials of influenza vaccine, 2 doses given in first year of life contain nearly 50 micrograms of mercury.
- Polysorbate 80: An emulsifier that crosses the blood brain barrier allowing vaccine ingredients into the brain. Causes infertility in animal studies.

 Human DNA: Many live viral vaccines are contaminated with fetal DNA debris & associated with autistic disorder and epidemic childhood leukemia and lymphomas. (Ref: T. Deisher).

The highest numbers of vaccine doses are given in the first 2 years of life during critical phases of brain and immune system development. Vaccines can trigger brain inflammation, derail normal brain development, impair thinking, concentration, attention, behavior, language and cause neurological motor injuries. Vaccines are not evaluated for carcinogenic or mutagenic potentials or impairment of fertility. Read Product Monographs available at www.vaccines411.ca under the health professional tab.

Make an informed vaccine decision!

Your Child. Your Future. Your Choice. www.vaccinechoicecanada.com

